



06-02-2023  
4th Copy

Change in pack size  
60 to 90ml

## Bisleri<sup>®</sup> Chewable Tablets / Capsules

(Iron (III) Hydroxide Polymaltose Complex + Folic Acid)

## Bisleri<sup>®</sup> Syrup

(Iron (III) Hydroxide Polymaltose Complex)

### DESCRIPTION

**Bisleri<sup>®</sup>** is a novel iron preparation used in the treatment of iron deficiency anemia. Iron, an essential constituent of the body, is necessary for haemoglobin formation and for the oxidative process of living tissue.

**Bisleri<sup>®</sup>** (Iron (III) Hydroxide Polymaltose Complex) is a low molecular weight carbohydrate polymer (Polymaltose). This prevents the iron from causing any harm in the gastrointestinal system. This protection inhibits interactions of the iron with food. Moreover, it ensures the bioavailability of the iron.

### COMPOSITION

#### **Bisleri<sup>®</sup> Chewable Tablets**

Each chewable tablet contains:  
Iron (III) Hydroxide Polymaltose Complex MS  
equivalent to Elemental Iron .....100mg  
Folic Acid USP.....350mcg

#### **Bisleri<sup>®</sup> Capsules**

Each capsule contains:  
Iron (III) Hydroxide Polymaltose Complex MS  
equivalent to Elemental Iron .....100mg  
Folic Acid USP.....350mcg

#### **Bisleri<sup>®</sup> Syrup**

Each 5ml contains:  
Iron (III) Hydroxide Polymaltose Complex MS  
equivalent to Elemental Iron .....50mg

### PROPERTIES AND ACTIONS

Iron Polymaltose, the active iron ingredient of essentially non-ionic, therefore, it has the following properties unlike ionized iron salt preparations:

- It does not cause irritation of the intestinal mucosa.
- It has palatable, non metallic taste (vanilla and toffee flavor).
- It has excellent tolerance.

**Bisleri<sup>®</sup>** can be taken with meals as appropriate to the patient, with any beverage or medications.

**Bisleri<sup>®</sup>** chewable tablets/capsules a novel preparation for the treatment of iron and folic acid deficiencies of all origins during pregnancy and lactation period. The folic acid content corresponds to the recommendation of the WHO.

**Bisleri<sup>®</sup>** syrup is pleasant tasted preparation tailored particularly for infants, children and elderly to cover the required demand of iron especially during pregnancy and lactation adolescents and woman capable of bearing children and adults.

### INDICATIONS:

#### **Bisleri<sup>®</sup> Chewable Tablets and Capsules:**

Prevention and treatment of iron and folic acid deficiencies of all origins during pregnancy and lactation period.

#### **Bisleri<sup>®</sup> Syrup:**

For infants, children and adults specifically women of child bearing age facing deficiency of iron.

### DOSAGE AND ADMINISTRATION:

Dosage and duration of therapy are dependent upon the extent of iron deficiency.

#### **Bisleri<sup>®</sup> Chewable Tablets and Capsules:**

One chewable tablet or capsule 2 or 3 times daily until a normalisation of haemoglobin value is achieved. Afterwards the therapy should be continued with one chewable tablet or capsule daily for a further month after normalisation of the blood values.

**Bisleri<sup>®</sup>** chewable tablets or capsules should be chewed or swallowed during or immediately after meal.

#### **Bisleri<sup>®</sup> Syrup:**

Dosage and duration of therapy are dependent upon the extent of iron deficiency and should be taken as directed by the physician.

Age Group	Manifest Iron Deficiency	Latent Iron Deficiency	Prophylactic Therapy
Infant (upto 1 year)	2.5 – 5ml daily	–	–
Children (upto 1-12)	5 – 10ml daily	2.5 – 5ml daily	–
Children (> 12 yrs)	10 – 30ml daily	5 – 10ml daily	–
Adults & nursing women	10 – 30ml daily	5 – 10ml daily	–
Pregnant Women	20 – 30ml daily	10ml daily	5 – 10ml daily

### CONTRAINDICATIONS:

- Overloading of iron (haemochromatosis, chronic haemolysis).
- Disturbances in iron utilisation (lead anemia, sidero-achrestic anemia), thalassemia.
- Megaloblastic anemia due to Vitamin B<sub>12</sub> deficiency.

### PRECAUTIONS:

As with all oral iron preparations, a dark colouration of the stool may occur which is without clinical significance.

### UNDESIRABLE EFFECTS:

Occasionally GI irritation, pressure in the epigastric region, nausea, constipation or diarrhoea may occur. Allergic reactions have been reported rarely with folic acid.

### INTERACTIONS:

Since, the iron is complex bound, ionic interactions with foodstuff components (phytins, oxalates, tannin, etc.) and concomitantly administered medications (tetracycline, antacids) are unlikely to occur.

### SHELF LIFE

See expiry on the pack

### AVAILABILITY

**Bisleri<sup>®</sup>** chewable tablets in a pack of 30's

**Bisleri<sup>®</sup>** capsules in a pack of 30's

**Bisleri<sup>®</sup>** syrup in a pack of 90ml and 120ml

### INSTRUCTIONS

**Dosage:** As directed by the physician.

Keep out of reach of children.

**For Chewable Tablets / Capsules:** Avoid exposure to heat, light and humidity.

Store between 15 to 30°C.

**For Syrup:** Do not store over 30°C, and protect from heat, light and freezing.

Medicine should not be used if container is leaking or it contains undissolved particles(s).

Improper storage may deteriorate the medicine.

Manufactured by:  
**SAMI Pharmaceuticals (Pvt.) Ltd.**  
F-95, S.I.T.E., Karachi-Pakistan  
www.samipharmapack.com  
Mfg. Lic. No. 000072

2000006311

**بسلیری** چیمو ایٹیل ٹیبلٹ / کیپسول / سیرپ  
(آئرن (III) ہائیڈرو آکسائیڈ پولی مالٹوز کمپلیکس + فولک ایسڈ)

ہدایات:

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بچوں کی پہنچ سے دور رکھیں۔

بچے کے لیے استعمال کیے جانے والے دوا کو گرمی، روشنی اور نمی سے محفوظ

۱۵ سے ۳۰ ڈگری سینٹی گریڈ کے درمیان میں رکھیں۔

بچے کے لیے سیرپ: دوا کو ۳۰ ڈگری سینٹی گریڈ سے زیادہ درجہ حرارت پر نہ رکھیں،

گرمی، روشنی اور نم ہونے سے محفوظ رکھیں۔

دوا کے ٹیک ہونے یا اس میں کوئی غیر حل پذیر شے نظر آنے کی صورت میں ہرگز استعمال نہ کریں۔

ورنہ دوا خراب ہو جائے گی۔

R.N-11/NA/02/2023\_For Syrup.

210mm

150mm