

Conspic® Tablets / Drops

(Sodium Picosulfate)

COMPOSITION:

Conspic® 5mg Laxative tablets:

Each tablet contains:
Sodium Picosulfate BP.....5mg

Conspic® Laxative Drops:

Each ml contains:
Sodium Picosulfate BP.....7.5mg

DESCRIPTION:

Picosulfate is the active ingredient in **Conspic®** which is a stool softening laxative for oral use

PHARMACOKINETICS AND PHARMACODYNAMICS:

Sodium picosulfate is chemically bisodic salt of 4-4'(2-Picollicliden) – bis (-Phenylsulphuric acid). It acts as a local laxative, metabolized by the colonic bacteria to the active compound bis (P-hydroxyphenyl) pyridyl-2-methane, which stimulate the mucosa of the large intestine & causing colonic peristalsis. Sodium picosulfate is usually effective within 10–14 hours

INDICATIONS:

Sodium picosulfate treats constipation by enhancing bowel movement, it is indicated in:

- Acute and chronic constipation
- Relief of evacuation in painful conditions such as hemorrhoids
- Pre and post operatively
- Preparation of barium enema
- Preparation of colon for proctosigmoidoscopy

DOSAGE AND ADMINISTRATION:

Adults:

1 - 2 tablets or 10 - 15 drops once a day

Children:

2 - 5 Years: 2 - 3 drops once daily

6 - 10 Years: 2 - 5 drops once daily

The dosage may be adjusted under medical advice, according to the age and symptoms
Drink 6 - 8 glasses of water daily while using any laxative

OR

As directed by the physician

CONTRAINDICATIONS:

Conspic® is contraindicated in patients with ileum, intestinal obstruction, acute abdominal conditions like appendicitis, acute inflammatory bowel disease and in severe dehydration. Known hypersensitivity to sodium picosulfate

PRECAUTIONS:

Prolonged use of **Conspic®** should be avoided. Prolong use may lead to diarrhoea, excessive loss & imbalance of electrolytes with possible hypokalemia. Children should not take **Conspic®** without medical advice. Do not use this laxative for longer than 1 week unless approved by doctor. Laxative may be habit-forming and can harm bowels if used too long

ADVERSE REACTIONS:

Following side effects related to the gastrointestinal tract may rarely be seen with sodium picosulfate use: abdominal pain, abdominal discomfort or distension, borborygmus, nausea, vomiting and diarrhoea

PREGNANCY AND LACTATION:

The safety of sodium picosulfate in pregnant and lactating women has not been established. Therefore sodium picosulfate should be administered to pregnant women only when expected therapeutic benefits are considered to outweigh any possible risks associated with treatment

STABILITY:

See expiry on the pack

PRESENTATION:

Conspic® 5mg laxative tablets in a pack of 30's

Conspic® laxative drops in a pack of 15ml

INSTRUCTION:

Keep out of reach of children
Avoid exposure to heat, light, humidity and freezing
Store between 15 to 30°C
Improper storage may deteriorate the medicine

Caution: Medicine should not be used if container is leaking
or it contains undissolved particle(s)

کونسیک ٹیبلٹ / ڈراپس
(سڈیم پیکوسلفیٹ)

خوراک: ڈاکٹری ہدایت کے مطابق استعمال کریں

بچوں کی پہنچ سے دور رکھیں

دوا کو صوب، گرمی، نمی اور نمند ہونے سے محفوظ 15 سے 30 ڈگری سینٹی گریڈ

کے درمیان میں رکھیں ورنہ دوا خراب ہو جائیگی

صحیح: دوا کے ٹک ہونے، دھملا ہونے یا اس میں کوئی غیر حل ہندے

ظہر آئے کی صورت میں ہرگز استعمال نہ کریں



Manufactured by:
SAMI Pharmaceuticals (Pvt) Ltd.
F-95, S.I.T.E., Karachi-Pakistan
www.samipharmap.com

P002142/S

R.N-01/HA/08/16