



10-05-2023
3rd Copy

(Change of Pack Size)

210mm

Dicloran[®] Gel

(Diclofenac Diethyl Ammonium Salt)

COMPOSITION:

Dicloran[®] Gel:

Each 100g contains:
Diclofenac Diethyl Ammonium Salt BP
equivalent to Diclofenac Sodium 2g

DESCRIPTION:

Dicloran[®] (Diclofenac) is a non steroidal anti-inflammatory drug (NSAID). It inhibits prostaglandins synthesis by decreasing the activity of the enzyme, cyclo-oxygenase, which results in decreased formation of prostaglandin precursors.

PHARMACODYNAMICS:

Dicloran[®] Gel relieves pain, reduce edema and shorten the time to return to normal function in inflammation of traumatic or rheumatic origin.

PHARMACOKINETICS:

Absorption:

The amount of Diclofenac absorbed through the skin is proportional to the contact time and skin area covered with **Dicloran[®] Gel**, topical dose and hydration of skin.

Distribution:

After topical administration to hand and knee joints, Diclofenac can be measured in plasma, synovial tissue, and synovial fluid.

Elimination:

Diclofenac and its metabolites are mainly excreted in urine.

INDICATIONS:

Post-traumatic inflammation of the tendons, ligaments, muscles, and joints, e.g. due to sprains, strains, and bruises. Localised form of soft-tissue rheumatism, e.g. tendovaginitis, bursitis, shoulder-hand syndrome and periarthropathy. Localised form of degenerative rheumatism, e.g. osteoarthritis of the peripheral joints and of the vertebral column.

DOSAGE AND ADMINISTRATION:

Apply 1 - 2g Gel 3 to 4 times daily on the affected area and rub gently. After application the hands should be washed, unless they are the site being treated. The duration of treatment depends on the indication and the response obtained. It is recommended that treatment be reviewed after 2 weeks Or as directed by the physician.

CONTRAINDICATIONS:

Known hypersensitivity to Diclofenac. It is also contraindicated in patients in whom attacks of asthma, urticaria or acute rhinitis are precipitated by acetylsalicylic acid or other NSAIDs.

PRECAUTIONS:

It should be applied only to healthy and intact skin surfaces. It should not be allowed to come in contact with the eyes or with mucous membrane. Not to be taken with mouth. Should be used for topical application only.

PREGNANCY:

Not recommended for use during pregnancy.

ADVERSE REACTIONS:

Occasionally allergic or non-allergic contact dermatitis. In isolated cases generalized skin rashes, hypersensitivity reactions, photosensitivity reactions may occur.

SHELF LIFE:

See expiry on the pack.

AVAILABILITY:

Dicloran[®] Gel in a tube of 30g.

INSTRUCTIONS:

For external use only

Keep out of reach of children

Do not store over 30°C, and protect from heat, light and freezing

Improper storage may deteriorate the medicine

ڈیکلوران جیل
(ڈیکلو فینیک ڈائی ایٹھائل امونیوم سالٹ)

ہدایات:

صرف بیرونی استعمال کے لئے ہے

بچوں کی پہنچ سے دور رکھیں

دوا کو 30°C سے زیادہ درجہ حرارت پر نہ رکھیں،

گرمی، روشنی اور نمند ہونے سے محفوظ رکھیں ورنہ دوا خراب ہو جائیگی

Manufactured by:
SAMI Pharmaceuticals (Pvt.) Ltd.
F-95, Off Hub River Road, S.I.T.E., Karachi-Pakistan
www.samipharmapk.com
Mfg. Lic. No. 000072

2000006660

R.N-03/NA/05/2023

120mm