


## 15-03-2022 1st Copy

	ости с
U	OCTILE <sup>3</sup> g Sachet
(Dio	ctahedral Smectite)
	TATIVE AND QUANTITATIVE COMPOSITION
	ILE <sup>®</sup> 3g Sachet
	shet contains:
Dioctane	dral Smectite MS3g
	IACEUTICAL FORM or oral suspension, Sachet.
	AL PARTICULARS
	EUTIC INDICATIONS:
	tomatic treatment of irritable bowel syndrome in adults and gastroesophageal reflux in infants due to mucoprotective effects. ment of acute and chronic diarrhoea.
POSOL	DGY AND METHOD OF ADMINISTRATION:
Children	
	the age of 1 year: 1 sachet per day.
	1 to 2 years: 1 to 2 sachets per day. the age of 2 years: 2 to 3 sachets per day.
	ministered in 2-3 divided doses.
	OR
	As prescribed by the physician
The cont	ents of the sachet can be mixed in a bottle with 50ml of water or with a semi-liquid food: broth, fruit, mashed vegetables, baby food.
Adults:	
An avera	ge of 3 sachets in 2-3 divided doses per day mixed in half a glass of water.
In acute	diarrhoea, the daily dosage can be doubled at the beginning of treatment.
	OR
	As prescribed by the physician
Administ	er preferably after meals in oesophagitis and in between meals in other indications.
CONTR	AINDICATIONS:
<ul> <li>Hyper</li> </ul>	rsensitivity to the active substance.
<ul> <li>Due to</li> </ul>	o the presence of glucose and saccharose, this medication is contraindicated for patients who are fructose intolerant.
	WARNINGS AND PRECAUTIONS FOR USE:
Treatmer	t with dioctahedral smectite does not eliminate the need of rehydration when it is required.
The volu	me of rehydration and its route of administration (Oral or IV) should be adapted to the patient's age and condition as well as to the severity of dia
	CTION WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION:
	dsorbent properties of this product may interfere with the rates and/or levels of absorption of other substances, it is recommended not to administer an ne same time as dioctahedral smectite.
-	
	TY, PREGNANCY AND LACTATION: , no malformation or foetotoxic effect has currently come to light. However, there is not enough evidence of pregnancies exposed to dioctahedral sme
exclude a	any risk.
Consequ	ently, given that dioctahedral smectite is not absorbed, its use could be envisaged during pregnancy only if necessary.
	S ON ABILITY TO DRIVE AND USE MACHINES:
Not know	m.
UNDESI	RABLE EFFECTS:
Very rare	cases of development or aggravation of constipation are known to be reported; treatment may be continued after decreasing the dosage.
	DSE:
OVERDU	
None	IACOLOGICAL PROPERTIES
None	IACOLOGICAL FROFLICTLS
None PHARN	cotherapeutic group: other intestinal absorbents. ATC Code: A07BC05.
None PHARM Pharmad	cotherapeutic group: other intestinal absorbents. ATC Code: A07BC05.
None PHARM Pharmac PHARM/	cotherapeutic group: other intestinal absorbents. ATC Code: A078C05. ACODYNAMIC PROPERTIES: dral smectile is a double aluminum and magnesium silicate; due to its structure in leaves and its high plastic viscosity, has a significant digestive n

— 115mm –





115mm