

# MOVERYL® Capsules (Celecoxib)

## COMPOSITION:

### Moveryl® 100mg Capsules:

Each capsule contains: Celecoxib MS..... 100mg

### Moveryl® 200mg Capsules:

Each capsule contains: Celecoxib MS..... 200mg

## PHARMACOLOGY:

### Pharmacodynamics:

Moveryl® (Celecoxib) is a non steroidal anti-inflammatory drug that exhibits anti-inflammatory, analgesic and anti-pyretic activities. The mechanism of action of Celecoxib is believed to be due to inhibition of prostaglandin synthesis, primarily via inhibition of cyclo-oxygenase-2 (Cox-2)

### Pharmacokinetics:

Peak plasma levels of Celecoxib occur approximately 3 hours after an oral dose. In healthy subjects, Celecoxib is highly protein bound (~97%) within the clinical dose range. Its metabolism is primarily mediated via cytochrome P450 2C9 and excretion is predominantly in bile with little unchanged drug recovered in urine and feces

## INDICATIONS AND USAGE:

Moveryl® (Celecoxib) is indicated for relief of the signs and symptoms of osteoarthritis and rheumatoid arthritis, for the treatment of primary dysmenorrhoea and for the management of acute pain, such as post-operative and dental pain

## DOSAGE AND ADMINISTRATION:

**Osteoarthritis:** Recommended dose is 200mg/day as a single dose or as 100mg twice per day. **Rheumatoid arthritis:** 100 to 200mg twice per day. **Management of acute pain and treatment of dysmenorrhoea:** 400mg initially, followed by an additional dose of 200mg, if needed on the first day. On subsequent days 200mg twice daily as needed. No dosage adjustment is necessary in case of elderly or in-patients with mild hepatic or renal impairment

OR

as directed by the physician

## CONTRA-INDICATIONS:

Known hypersensitivity to the drug or other sulfonamides. It is also contra-indicated in patients who have experienced asthma, urticaria or allergic-type reactions after taking aspirin or other NSAIDs

## WARNINGS AND PRECAUTIONS:

Celecoxib should not be used in patients having ulcer, asthma, urticaria or acute rhinitis. No clinical experiences are available in patients with severe hepatic and renal impairment. Safety of the drug during pregnancy and lactation and in children below 18 years has not been established

## DRUG INTERACTIONS:

Celecoxib metabolism is predominantly mediated via cytochrome P 450 2C9 in the liver. Co-administration of it with the drugs that are known to inhibit 2C9 should be done with caution

### ACE inhibitors and Diuretics:

Concomitant use with ACE inhibitors or thiazides may reduce the antihypertensive or diuretic effects respectively

### Aspirin:

Co-administration with aspirin increases the rate of GI ulceration or other complications

### Fluconazole:

Fluconazole increases two fold the plasma concentration of Celecoxib

## ADVERSE EFFECTS:

Headache, sinusitis, backache, abdominal pain, diarrhoea, dyspepsia, nausea, rash, insomnia etc. Rarely gastro-intestinal bleeding and anaphylactic reaction may occur

## OVER DOSAGE:

Patients should be managed by symptomatic and supportive care following an over dose. There are no specific antidotes

## STABILITY:

See expiry on the pack

## PRESENTATIONS:

Moveryl® 100mg Capsules in pack of 20's

Moveryl® 200mg Capsules in pack of 20's

## INSTRUCTIONS:

Keep out of reach of children

Avoid exposure to heat, light and humidity

Store at 30°C or below

Improper storage may deteriorate the medicine

موورل کیپسول  
(سیلیکوکسیب)

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں

ہدایات: بچوں کی پہنچ سے دور رکھیں

دوا کو دھوپ، گرمی اور نمی سے محفوظ ۳۰ ڈگری سینٹی گریڈ

تھلا سے کم درجہ حرارت پر رکھیں ورنہ دوا خراب ہو جائے گی



Manufactured by:  
**SAMI PHARMACEUTICALS (PVT) LTD.**  
F-95, S.I.T.E., Karachi-Pakistan  
www.samipharmapk.com