



09-04-2022
1st Copy

Change of Pack Size

210mm

Nims[®] Tablets (Nimesulide)

COMPOSITION

Nims[®] 100mg Tablets

Each film coated tablet contains:
Nimesulide Ph. Eur.....100mg

PHARMACOLOGY

Pharmacodynamics:

Nimesulide is a non-steroidal anti-inflammatory drug that selectively inhibits cyclooxygenase-2 but it appears to exert its therapeutic effects through variety of other mechanisms as well like inhibition of platelet activating factor (PAF), tumor necrosis factor-alpha (TNF- α), protease & histamine etc.

Pharmacokinetics:

Oral absorption of Nimesulide is almost complete. Concomitant administration of food may decrease the rate but not the extent of absorption of Nimesulide. Plasma binding is high (approximately 99%). Nimesulide is extensively metabolized to several metabolites which are excreted mainly in the urine (about 70%) and in feces (about 20%)

USAGE

Nimesulide, being a NSAID, exhibits anti-inflammatory & analgesic properties. It is indicated as second choice for the treatment of acute pain and primary dysmenorrhoea.

SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Hepatic Effects:

Rarely Nimesulide has been reported to be associated with serious hepatic reactions, including very rare fatal cases. Patients who experience symptoms compatible with hepatic injury during treatment with Nimesulide (e.g. anorexia, nausea, vomiting, abdominal pain, fatigue, dark urine) or patients who develop abnormal liver function tests should have treatment discontinued. These patients should not be rechallenged with Nimesulide. Liver damage, in most cases reversible, has been reported following short exposure to the drug

CONTRAINDICATIONS

The use of Nimesulide is contraindicated in patients with hepatic impairment, children, active peptic ulcer disease, known hypersensitivity to the active substance, acetylsalicylic acid, or other prostaglandin synthetase inhibiting drugs

ADVERSE REACTIONS

Undesirable effects include gastrointestinal disorders, gastrointestinal bleeding, duodenal ulcers and perforation, gastric ulcer and perforation

SIDE-EFFECTS

At therapeutic doses Nimesulide is usually well tolerated. Occasional side effects include heartburn, nausea and epigastric pain. Skin rashes of allergic type may rarely occur. Uncommon side effects include bleeding from stomach or bowel; duodenal or stomach ulcer and burst ulcers

DRUG INTERACTIONS

Clinically important interactions involving interference with drug metabolism have not been described with Nimesulide

DOSAGE AND ADMINISTRATION

Usual oral dose in adults is one tablet twice daily for not more than 15 days

OR

As directed by the physician

SHELF LIFE

See expiry on the pack

AVAILABILITY

Nims[®] 100mg tablets in a pack of 30's

INSTRUCTIONS

Keep out of reach of children

Avoid exposure to heat, light and humidity

Store between 15 to 30°C

Improper storage may deteriorate the medicine

نیمز[®]
ٹیبلٹ
(نیمی سولائیڈ)

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں۔

بچوں کی پہنچ سے دور رکھیں۔

دوا کو گرمی، روشنی اور نمی سے محفوظ ۱۵ سے ۳۰ ڈگری سینٹی گریڈ

کے درمیان میں رکھیں ورنہ دوا خراب ہو جائیگی۔

Manufactured by:
SAMI Pharmaceuticals (Pvt.) Ltd.
F-95, S.I.T.E., Karachi-Pakistan
www.samipharmapk.com
Mfg. Lic. No. 000072

2000005508

R.N-06/NA/04/2022

120mm