

Pulsate 100mg Tablets

Pulsate® 100mg Dispersible Tablets
(Roxithromycin)

COMPOSITION:

Each dispersible tablets contains:
Roxithromycin EP100mg

INDICATIONS:

PULSATE® is an antibiotic of the macrolide group. It is recommended for ENT, bronchopulmonary, skin infections and in the prevention of meningococcal infections

CONTRA-INDICATIONS:

PULSATE® must not be used in the following cases:

- Known allergy to macrolides, such as erythromycin, josamycin etc.
- In combination with ergot derivatives

Possible side-effects:

Gastro-intestinal reactions: Nausea, vomiting, stomach pain, diarrhoea, Allergic skin reactions, Dizzy spells

DOSAGE:

The dosage to be used is 5 to 8mg/kg/day in 2 separate doses and treatment should not prolong for more than 10 days

- Children between 24 and 40 kgs: 100mg morning and evening
- The tablets should preferably be taken before meals

ADMINISTRATION:

Disperse the tablet in 2 to 3 teaspoonsful freshly boiled & cooled water before use

PRECAUTIONS:

Pulsate® is not recommended in Hepatic insufficiency, in combination with ergot derivatives, particularly ergotamine

PRESENTATION:

Pulsate® 100mg Tablets in pack of 10's

STABILITY:

See expiry on the pack

INSTRUCTIONS:

Keep out of reach of children
Avoid exposure to heat, light and humidity
Store at 25°C or below
Improper storage may deteriorate the medicine

Manufactured by:
SAMI Pharmaceuticals (Pvt) Limited
F-95, S.I.T.E., Karachi-Pakistan

پلسیٹ ۱۰۰ ملی گرام ڈیسپرسبل ٹیبلٹ
(روکسی تھرو مائین)
خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں
ہدایت: بچوں کی پہنچ سے دور رکھیں
دوا کو صاف، گرمی اور نمی سے محفوظ ۲۵ ڈگری سینٹی گریڈ
یا اس سے کم درجہ حرارت پر رکھیں ورنہ دوا اثر ناپاؤں ہو جائے گی

Pulsate 150mg & 300mg Tablets

Pulsate® Tablets (Roxithromycin)

Composition:

Pulsate® 150mg tablet contains: Roxithromycin.....150mg

Pulsate® 300mg tablet contains: Roxithromycin.....300mg

INDICATIONS:

Pulsate® is an antibiotic of the macrolide group. It is recommended for ENT, bronchopulmonary, genital (excluding gonococcal infections) and skin manifestations

CONTRA-INDICATIONS:

Pulsate® must not be used in following cases:

- Allergy to macrolides, such as roxithromycin, erythromycin, josamycin etc.
- In combination with ergotamine and dihydroergotamine
- 300mg tablet is contra-indicated in patients with severe hepatic insufficiency

PRECAUTIONS:

The physician should be informed in the case of:

- Hepatic insufficiency
- Concomitant administration of other drugs
- Lactation
- Pregnancy

INTERACTIONS WITH OTHER SIDE-EFFECTS:

- There is no clinically significant interactions with warfarin carbamazepine, cyclosporin, oral contraceptives containing oestrogens and progestogens
- In healthy volunteers, a slight increase in plasma concentrations of theophylline levels which does not necessitate to alter the usual dosage, has been shown

SIDE EFFECTS:

Gastro-intestinal reactions: Gastric pain, nausea, vomiting, stomach pain, diarrhoea

Allergic skin reactions

Dizzy sensations

Moderate increase in ASAT, ALAT and/or alkaline phosphatases. Exceptionally acute cholestatic hepatitis

Possibility of fungal overgrowth

DOSAGE AND ADMINISTRATION:

Adults:

Pulsate® 150mg: 1 tablet twice daily preferably before meals

Pulsate® 300mg: 1 tablet daily preferably before meals

PRESENTATION:

Pulsate® 150mg tablets in pack of 10's

Pulsate® 300mg tablets in pack of 10's

STABILITY:

See expiry on the pack

INSTRUCTIONS:

Keep out of reach of children

Avoid exposure to heat, light and humidity

Store at 25°C or below

Improper storage may deteriorate the medicine

پلسیٹ ٹیبلٹ
(روکسی تھرو مائین)

خوراک: ڈاکٹر کی ہدایت کے مطابق استعمال کریں

ہدایت: بچوں کی پہنچ سے دور رکھیں

دوا کو چھپ، گرمی اور نمی سے محفوظ رکھیں 25°C یا اس سے کم درجہ حرارت پر رکھیں اور نندو اور خراب ہو جائے گی

Manufactured by:
SAMI Pharmaceuticals (Pvt) Limited
F-95, S.I.T.E., Karachi-Pakistan

